

MAT0028 ~ Lesson 33

Work the following examples as you listen to the recorded lecture.

Factoring Complex Trinomials

Problem type: $ax^2 + bx + c$ (Where a , b and c are numbers, and x is the unknown.)

Rules for signs: Use the same sign rules for all trinomials

Rule 1: If the **2nd sign is +**, then both factor signs will match the **1st sign** in the problem.

$$ax^2 + bx + c \rightarrow (\quad + \quad)(\quad + \quad) \quad ax^2 - bx + c \rightarrow (\quad - \quad)(\quad - \quad)$$

Rule 2: If the **2nd sign is -**, then the factor signs will be different, + and -.

$$ax^2 + bx - c \rightarrow (\quad + \quad)(\quad - \quad) \quad ax^2 - bx - c \rightarrow (\quad + \quad)(\quad - \quad)$$

Rule 3: Use 2nd operation to find out if you add or subtract factors to equal b .

Steps to remember:

1. Set the factor statement
2. Set the binomial factors with the signs
3. Factor the variable squares
4. Find all factor combinations for "a" and "c" until one matches the factor statement
5. Place the "a" factors in the 1st positions and the "c" factors in the 2nd positions of the binomial pairs
6. FOIL to check

Example 1: $25x^2 - 20x + 4$	Step 1: Factor Statement:
() ()	Step 2: Set signs for the factors.
	Step 3: Factor the variable squares.
	Step 4: Factor a; factor and reverse factor c $a = \underline{\hspace{2cm}}$ $c = \underline{\hspace{2cm}}$ <u>Factor Statement Work Space</u> _____ _____ _____ _____ _____ _____
	Look for the combination that fits the factor statement.
	Step 5: Use the "a" factors in the 1 st positions and the "c" factors in 2 nd positions of your solution.
	Step 6: FOIL to check.